

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

ALDH2 monoclonal antibody (M09), clone 2A7

Catalog Number: H00000217-M09

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against a partial recombinant ALDH2.

Clone Name: 2A7

 $\label{eq:mmunogen: ALDH2 (AAH02967, 408 a.a. ~ 517 a.a)} partial recombinant protein with GST tag. MW of the$

GST tag alone is 26 KDa.

Sequence:

DGMTIAKEEIFGPVMQILKFKTIEEVVGRANNSTYGLAA AVFTKDLDKANYLSQALQAGTVWVNCYDVFGAQSPF GGYKMSGSGRELGEYGLQAYTEVKTVTVKVPQKNS

Host: Mouse

Reactivity: Human

Applications: ELISA, S-ELISA, WB-Re

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 217

Gene Symbol: ALDH2

Gene Alias: ALDH-E2, ALDHI, ALDM, MGC1806

Gene Summary: This protein belongs to the aldehyde dehydrogenase family of proteins. Aldehyde dehydrogenase is the second enzyme of the major oxidative pathway of alcohol metabolism. Two major

isoforms of this enzyme, cytosolic and mitochondrial, be distinguished can by their electrophoretic mobilities, kinetic properties, and subcellular localizations. Most Caucasians have two major isozymes, while approximately 50% of Orientals only the cytosolic isozyme, mitochondrial isozyme. A remarkably higher frequency of acute alcohol intoxication among Orientals than among Caucasians could be related to the absence of the mitochondrial isozyme. This gene encodes mitochondrial isoform, which has a low Km acetaldehydes, and is localized in mitochondrial matrix. [provided by RefSeq]