

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## Datasheet

## CSE1L monoclonal antibody (M05), clone 3F8

Catalog Number: H00001434-M05

Regulation Status: For research use only (RUO)

**Product Description:** Mouse monoclonal antibody raised against a partial recombinant CSE1L.

Clone Name: 3F8

**Immunogen:** CSE1L (NP\_001307, 872 a.a. ~ 971 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

## Sequence:

LIGLFELPEDDTIPDEEHFIDIEDTPGYQTAFSQLAFAGK KEHDPVGQMVNNPKIHLAQSLHKLSTACPGRVPSMVS TSLNAEALQYLQGYLQAARVTLL

Host: Mouse

Reactivity: Human

Applications: ELISA, IF, IHC-P, S-ELISA, WB-Ce, WB-Re (See our web site product page for detailed application

(See our web site product page for detailed applications information)

**Protocols:** See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Isotype: IgG1 kappa

Storage Buffer: In 1x PBS, pH 7.4

**Storage Instruction:** Store at -20 °C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 1434

Gene Symbol: CSE1L

Gene Alias: CAS, CSE1, MGC117283, MGC130036, MGC130037, XPO2

**Gene Summary:** Proteins that carry a nuclear localization signal (NLS) are transported into the nucleus

by the importin-alpha/beta heterodimer. Importin-alpha binds the NLS, while importin-beta mediates translocation through the nuclear pore complex. After translocation, RanGTP binds importin-beta and displaces importin-alpha. Importin-alpha must then be returned to the cytoplasm, leaving the NLS protein behind. The protein encoded by this gene binds strongly to NLS-free importin-alpha, and this binding is released in the cytoplasm by the combined action of RANBP1 and RANGAP1. In addition, the encoded protein may play a role both in apoptosis and in cell proliferation. [provided by RefSeq]