

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

ZAP70 monoclonal antibody, clone SBZAP (AP)

Catalog Number: MAB5889

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against partial recombinant ZAP70.

Clone Name: SBZAP

Immunogen: Recombinant protein corresponding to

amino acids 280-309 of human ZAP70.

Host: Mouse

Reactivity: Human

Applications: ELISA, Flow Cyt

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Specificity: human TCR ?; Zap-70 (Mr 70 KDa).

Form: Liquid

Conjugation: AP

Concentration: Lot specific

Isotype: IgG1, kappa

Recommend Usage: ELISA (1:500-1:1000)

The optimal working dilution should be determined by

the end user.

Storage Buffer: In 50 mM Tris, pH 8.0 (1mM MgCl₂,

50% glycerol, 0.09% sodium azide)

Storage Instruction: Store in the dark at 4°C. Do not

freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 7535

Gene Symbol: ZAP70

Gene Alias: FLJ17670, FLJ17679, SRK, STD, TZK,

ZAP-70

Gene Summary: This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, combined а severe immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq]

References:

1. The Syk/ZAP-70 protein tyrosine kinase connection to antigen receptor signalling processes. van Oers NS, Weiss A. Semin Immunol. 1995 Aug;7(4):227-36.

2. F2(Pmp)2-TAM zeta 3, a novel competitive inhibitor of the binding of ZAP-70 to the T cell antigen receptor, blocks early T cell signaling. Wange RL, Isakov N, Burke TR Jr, Otaka A, Roller PP, Watts JD, Aebersold R, Samelson LE. J Biol Chem. 1995 Jan 13;270(2):944-8.