

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

## **Datasheet**

## CD99 monoclonal antibody (M01), clone 3A10

Catalog Number: H00004267-M01

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody

raised against a partial recombinant CD99.

Clone Name: 3A10

Immunogen: CD99 (AAH03147, 23 a.a. ~ 122 a.a) partial recombinant protein with GST tag. MW of the

GST tag alone is 26 KDa.

## Sequence:

DGGFDLSDALPDNENKKPTAIPKKPSAGDDFDLGDAV VDGENDDPRPPNPPKPMPNPNHPSSSGSFSDADL ADGVSGGEGKGGSDGGGSHRKEGEEAD

Host: Mouse

Reactivity: Human

**Applications:** ELISA, S-ELISA, WB-Ce, WB-Re, WB-Tr (See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG2a Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 4267

Gene Symbol: CD99

Gene Alias: MIC2, MIC2X, MIC2Y

**Gene Summary:** The protein encoded by this gene is a cell surface glycoprotein involved in leukocyte migration, T-cell adhesion, ganglioside GM1 and transmembrane protein transport, and T-cell death by a

caspase-independent pathway. In addition, the encoded protein may have the ability to rearrange the actin cytoskeleton and may also act as an oncosuppressor in osteosarcoma. Cyclophilin A binds to CD99 and may act as a signaling regulator of CD99. This gene is found in the pseudoautosomal region of chromosomes X and Y and escapes X-chromosome inactivation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]