pBABEhygro-RhoA L63 Retroviral Vector

CATALOG NUMBER: RTV-204 **STORAGE:** -80ºC

QUANTITY AND CONCENTRATION: 100 µL of bacterial glycerol stock

Background

Retroviruses are efficient tools for delivering heritable genes into the genome of dividing cells. Cell Biolabs' retrovirus vector is based on the pBABE vector system, which is derived from Moloney murine leukemia virus (MMLV). The vector provides the viral package signal, transcription and processing elements, and a target gene. The viral *env* gene, produced by the package cell line, encodes the envelop protein, which determines the viral infectivity range. Transfection into a package cell line produces high-titer, replication-incompetent viruses. In addition to transfer and expression of exogenous genes in mammalian cells, recently, retroviruses have been used to express silencing RNAs (siRNA) to decrease the expression of target genes both *in vitro* and *in vivo*.

The vector contains the bacterial origin of replication, ampicillin-resistance gene, and hygromycin-resistance gene for the growth of infected mammalian cells to select stable cell lines (Figure 1).

Three members of the Rho family of small GTPases, Rho, Rac, and Cdc42, have been shown to play a crucial role in regulating the organization of the actin cytoskeleton in response to extracellular stimuli. Activation of Rho, Rac, and Cdc42 in quiescent Swiss 3T3 fibroblasts induces the assembly of filamentous actin into stress fibers, lamellipodia, and filopodia, respectively. In addition to these effects on the actin cytoskeleton, it has been shown that Rac and Cdc42 (and in some cells Rho) can activate JNK and p38 that leads to transcriptional activation. In fibroblast cells, Rho, Rac, and Cdc42 have each been implicated in cell cycle control. A constitutively active form of human RhoA L63 sequence is cloned into the retroviral vector pBABEhygro at the *BamHI* and *Sna*B I sites with a myc tag at its N-terminus.

Safety Consideration

Remember that you will be working with samples containing infectious virus. Follow the recommended NIH guidelines for all materials containing BSL-2 organisms. Always wear gloves, use filtered tips and work under a biosafety hood.

References

1. Morgenstern, J. P. and H Land. (1990) *Nuc. Acid Res.* 18, 3587-3596.
2. Coffin, J. M. and H. E. Varmus, *Retroviruses*, Cold Spring Harbor Press, NY.
3. Schuck S, Manninen A, Honsho M, Fullekrug J and Simons K. (2004) *Proc Natl Acad Sci U S A.* 101, 4912-4917.
4. Machesky L. M. and Hall A. (1996) *Trends Cell Biol.* 6:304-10.




Figure 1. Retroviral Vector Map

Warranty

These products are warranted to perform as described in their labeling and in Cell Biolabs literature when used in accordance with their instructions. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THIS EXPRESSED WARRANTY AND CELL BIOLABS DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS FOR PARTICULAR PURPOSE. CELL BIOLABS's sole obligation and purchaser's exclusive remedy for breach of this warranty shall be, at the option of CELL BIOLABS, to repair or replace the products. In no event shall CELL BIOLABS be liable for any proximate, incidental or consequential damages in connection with the products.

This product is for RESEARCH USE ONLY; not for use in diagnostic procedures.

Contact Information

Cell Biolabs, Inc.
7758 Arjons Drive
San Diego, CA 92126
Worldwide: +1 858-271-6500
USA Toll-Free: 1-888-CBL-0505
E-mail: tech@cellbiolabs.com
www.cellbiolabs.com

©2004-2008: Cell Biolabs, Inc. - All rights reserved. No part of these works may be reproduced in any form without permissions in writing.