

prosci-inc.com




HIGH PERFORMANCE ANTIBODIES ... AND MORE

ProSci Incorporated 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

CD11b Antibody [ICRF44] (APC)

CATALOG NUMBER: 76-095



Human peripheral blood lymphocytes were stained with APC ICRF44 with relevant isotype control in Red.

Specifications	
SPECIES REACTIVITY:	Human
TESTED APPLICATIONS:	FACS
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
SPECIFICITY:	The ICRF44 monoclonal antibody specifically reacts with the 165 kDa human adhesion glycoprotein CD11b, which forms, together with the 95 kDa CD18 (integrin beta2) a complex known as Mac-1.
HOST SPECIES:	Mouse
Properties	
PURIFICATION:	The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.
PHYSICAL STATE:	liquid
BUFFER:	Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.
CONCENTRATION:	5 uL (1 ug) / test
STORAGE CONDITIONS:	The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze.
CLONALITY:	Monoclonal
ISOTYPE:	Mouse IgG1, kappa
CONJUGATE:	APC
Additional Info	
ALTERNATE NAMES:	CR3A, MO1A, CD11B, MAC-1, MAC1A, SLEB6, ITGAM
OFFICIAL SYMBOL:	ITGAM
GENE ID:	3684
Background	
BACKGROUND:	The ICRF44 monoclonal antibody specifically reacts with the 165 kDa human adhesion glycoprotein CD11b, which forms, together with the 95 kDa CD18 (integrin beta2) a complex known as Mac-1. CD11b is expressed

	on the surface of activated lymphocytes, a subset of natural killer cells, granulocytes, and monocytes. It functions as a receptor in cell-cell and cell-matrix interactions and as a receptor for iC3b, ICAM-1, ICAM-2, and ICAM-3 intercellular adhesion molecules.
REFERENCES:	1) Barclay, A. N., Brown, M. H., Law, S. A. K. A., McKnight, A. J., Tomlinson, M. G., van der Merwe, P. A. (1997). The leucocyte antigen factsbook. Academic Press.
	2) Knapp W(1989) Leucocyte typing IV: white cell differentiation antigens. Oxford University Press, 1989.
	 Sotiriou, S. N., Orlova, V. V., Al-Fakhri, N., Ihanus, E., Economopoulou, M., Isermann, B., Chavakis, T. (2006). Lipoprotein (a) in atherosclerotic plaques recruits inflammatory cells through interaction with Mac-1 integrin. The FASEB journal, 20(3), 559-561.

FOR RESEARCH USE ONLY

December 13, 2016