

prosci-inc.com





HIGH PERFORMANCE ANTIBODIES ... AND MORE

ProSci Incorporated 12170 Flint Place Poway, CA 92064

Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

CD44 Antibody [IM7] (FITC)

CATALOG NUMBER: 76-415

Specifications	
SPECIES REACTIVITY:	Human, Mouse
TESTED APPLICATIONS:	FACS
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
SPECIFICITY:	The IM7 monoclonal antibody specifically reacts with all the isoforms and both alloantigens of the CD44 glycoprotein (Pgp-1, Ly-24).
HOST SPECIES:	Rat
Properties	
PURIFICATION:	The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.
PHYSICAL STATE:	liquid
BUFFER:	Phosphate-buffered aqueous solution, ≤0.09% Sodium azide, may contain carrier protein/stabilizer, ph7.2.
CONCENTRATION:	0.5 mg/mL
STORAGE CONDITIONS:	The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze.
CLONALITY:	Monoclonal
ISOTYPE:	Rat IgG2b, kappa
CONJUGATE:	FITC
Additional Info	
ALTERNATE NAMES:	IN, LHR, MC56, MDU2, MDU3, MIC4, Pgp1, CDW44, CSPG8, HCELL, HUTCH-I, ECMR-III, Ly-24, Pgp-1, HERMES, AU023126, AW121933, AW146109, CD44
OFFICIAL SYMBOL:	CD44
GENE ID:	960; 12505
Background	
BACKGROUND:	The IM7 monoclonal antibody specifically reacts with all the isoforms and both alloantigens of the CD44 glycoprotein (Pgp-1, Ly-24). CD44 is expressed on hematopoietic and non-hematopoietic cells, bone marrow myeloid cells, memory T lymphocytes, periphery activated B cells, CD4+ T lymphocytes, and CD8+ T lymphocytes. The periphery B and T lymphocytes upregulate the expression of CD44. CD44 binds to hyaluronar molecules, acting as an adhesion molecule. The IM7 antibody inhibits collagen-induced arthritis in DBA/1 mice, prevents central nervous system inflammation associated with experimental autoimmune encephalomyelitis, but exacerbates the experimental autoimmune thyroiditis in CBA/J mice. The IM7 antibody also cross-reacts with dog, human, pig, horse, cat, and cow leukocytes.
REFERENCES:	1) Trowbridge, I. S., Lesley, J., Schulte, R., Hyman, R., Trotter, J. (1982). Biochemical characterization and cellular distribution of a polymorphic, murine cell-surface glycoprotein expressed on lymphoid tissues.lmmunogenetics,15(3), 299-312.
	2) Lesley, J., Trowbridge, I. S. (1982). Genetic characterization of a polymorphic murine cell-surface

glycoprotein.Immunogenetics,15(3), 313-320.

3) Nedvetzki, S., Walmsley, M., Alpert, E., Williams, R. O., Feldmann, M., Naor, D. (1999). CD44 involvement in experimental collagen-induced arthritis (CIA). Journal of autoimmunity, 13(1), 39-47.

FOR RESEARCH USE ONLY

December 13, 2016