

Orders: orders@novusbio.com

Support: technical@novusbio.com

Web: www.novusbio.com

Protocols, Publications, Related Products, Reviews and more:

www.novusbio.com/NBP2-05316

NBP2-05316

TSGA10IP Overexpression Lysate

Specificity/Sensitivity Homo sapiens TSGA10IP, mRNA.

Immunogen

The lysate was created in HEK293T cells, using plasmid ID RC215316 and based on accession number NP_689975. The protein contains a C-terminal DDK tag.

Recommended Dilutions

Buffer

RIPA buffer

Uses

This product is intended for use as a positive control in Western Blot. You will receive the lysate (100ug), and an empty vector negative control (100 ug).

Images

Western Blot: TSGA10IP Overexpression Lysate (Adult Normal) [NBP2-05316] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for TSGA10IP.

Notes

This product is manufactured by and sold under license from OriGene Technologies and its use is limited solely for research purposes. Please note that '1 mg/ml' refers to the total protein concentration and that the specific protein concentration is not determined at this time.

Updated 3/9/2013 2.0

Limitations: This product is for research use only and is not approved for use in humans or in clinical diagnosis. Products are guaranteed for six months from the date of receipt except for peptides and proteins which are guaranteed for 3 months. For more information on our guarantee, please visit www.novusbio.com/guarantee.

Novus USA

8100 Southpark Way, A-8 Littleton, CO 80120 p) 888-506-6887 p) 303-730-1950 f) 303-730-1966 novus@novusbio.com

Novus Canada

461 North Service Road West, Unit B37 Oakville, ON L6M 2V5 p) 855-668-8722 f) 905-827-6402 canada@novusbio.com

Novus Europe

12 Cambridge Science Park Cambridge, UK CB4 0FQ UK: p) +44-1223-426-001, f) +44-871-971-1635 DE: p) +49-6922-22340-60, f) +49-0800-58926-79 IT: p) +39-02-4032-6786, f) +39-02-4032-6340 FR: p) +33-1-76-77-45-30, f) +33-1-76-77-45-31 europe@novusbio.com



olely for rese tration and th

TSGA1 Unit Size

Storage Store at -20C. Avoid freeze-thaw cycles.

Applications WB

Purity

Protein

Molecular Weight 62.2kDa