

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

ARHGDIA monoclonal antibody (M02), clone 1G5-2F3

Catalog Number: H00000396-M02

Regulation Status: For research use only (RUO)

Product Description: Mouse monoclonal antibody raised against a full length recombinant ARHGDIA.

Clone Name: 1G5-2F3

 $\label{lem:mmunogen: ARHGDIA (AAH16031, 1 a.a. \sim 204 a.a)} % \begin{subarray}{ll} \textbf{MW} & \textbf{MW} &$

GST tag alone is 26 KDa.

Sequence:

MAEQEPTAEQLAQIAAENEEDEHSVNYKPPAQKSIQEI QELDKDDESLRKYKEALLGRVAVSADPNVPNVVVTGL TLVCSSAPGPLELDLTGDLESFKKQSFVLKEGVEYRIKI SFRVNREIVSGMKYIQHTYRKGVKIDKTDYMVGSYGP RAEEYEFLTPVEEAPKGMLARGSYSIKSRFTDDDKTD HLSWEWNLTIKKDWKD

Host: Mouse

Reactivity: Human

Applications: ELISA, S-ELISA, WB-Re

(See our web site product page for detailed applications

information)

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Isotype: IgG1 Kappa

Storage Buffer: In 1x PBS, pH 7.4

Storage Instruction: Store at -20°C or lower. Aliquot to

avoid repeated freezing and thawing.

Entrez GenelD: 396

Gene Symbol: ARHGDIA

Gene Alias: GDIA1, MGC117248, RHOGDI, RHOGDI-1

Gene Summary: Aplysia Ras-related homologs (ARHs),

also called Rho genes, belong to the RAS gene superfamily encoding small guanine nucleotide exchange (GTP/GDP) factors. The ARH proteins may be kept in the inactive, GDP-bound state by interaction with GDP dissociation inhibitors, such as ARHGDIA (Leffers et al., 1993 [PubMed 8262133]).[supplied by OMIM]

References:

1. Comparative proteomic analysis on human L-02 liver cells treated with varying concentrations of trichloroethylene Liu J, Huang H, Xing X, Xi R, Zhuang Z, Yuan J, Yang F, Zhao J. Toxicol Ind Health. 2007 Mar;23(2):91-101.