

Datasheet

ZAP70 monoclonal antibody, clone SBZAP (Biotin)

ZAP-70

Catalog Number: MAB5891**Regulation Status:** For research use only (RUO)**Product Description:** Mouse monoclonal antibody raised against partial recombinant ZAP70.**Clone Name:** SBZAP**Immunogen:** Recombinant protein corresponding to amino acids 280-309 of human ZAP70.**Host:** Mouse**Reactivity:** Human**Applications:** ELISA, Flow Cyt
(See our web site product page for detailed applications information)**Protocols:** See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols**Specificity:** human TCR ?; Zap-70 (Mr 70 KDa).**Form:** Liquid**Conjugation:** Biotin**Isotype:** IgG1, kappa**Recommend Usage:** The optimal working dilution should be determined by the end user.**Storage Buffer:** In PBS (0.09% sodium azide)**Storage Instruction:** Store in the dark at 4°C. Do not freeze.

Avoid prolonged exposure to light.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 7535**Gene Symbol:** ZAP70**Gene Alias:** FLJ17670, FLJ17679, SRK, STD, TZK,

Gene Summary: This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq]

References:

1. The Syk/ZAP-70 protein tyrosine kinase connection to antigen receptor signalling processes. van Oers NS, Weiss A. Semin Immunol. 1995 Aug;7(4):227-36.
2. F2(Pmp)2-TAM zeta 3, a novel competitive inhibitor of the binding of ZAP-70 to the T cell antigen receptor, blocks early T cell signaling. Wange RL, Isakov N, Burke TR Jr, Otaka A, Roller PP, Watts JD, Aebersold R, Samelson LE. J Biol Chem. 1995 Jan 13;270(2):944-8.