

9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

Datasheet

Cd3e monoclonal antibody, clone 145-2C11 (PE)

Catalog Number: MAB5909

Regulation Status: For research use only (RUO)

Product Description: Hamster monoclonal antibody raised against Cd3e.

Clone Name: 145-2C11

Immunogen: H-2Kb-specific mouse cytotoxic T lymphocyte clone BM10-37.

Host: Hamster

Reactivity: Mouse

Applications: Flow Cyt, IP, WB (See our web site product page for detailed applications information)

Protocols: See our web site at http://www.abnova.com/support/protocols.asp or product page for detailed protocols

Specificity: Mouse CD3? (Mr 25 KDa).

Form: Liquid

Conjugation: PE

Isotype: IgG, kappa

Recommend Usage: Flow Cytometry $(0.1 \text{ ug}/10^6 \text{ cells})$ The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS (0.09% sodium azide)

Storage Instruction: Store in the dark at 4°C. Do not freeze. Avoid prolonged exposure to light. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 12501

Gene Symbol: Cd3e

Gene Alias: Al504783, CD3, CD3epsilon, T3e

References:

1. Differential activation of antigen-stimulated suicide and cytokine production pathways in CD4+ T cells is regulated by the antigen-presenting cell. Wang R, Murphy KM, Loh DY, Weaver C, Russell JH. J Immunol. 1993 May 1;150(9):3832-42.

2. Activation-driven T cell death. II. Quantitative

differences alone distinguish stimuli triggering nontransformed T cell proliferation or death. Ucker DS, Meyers J, Obermiller PS. J Immunol. 1992 Sep 1;149(5):1583-92.

3. Circulating CD3+/T cell receptor V gamma 3+ fetal murine thymocytes home to the skin and give rise to proliferating dendritic epidermal T cells. Payer E, Elbe A, Stingl G. J Immunol. 1991 Apr 15;146(8):2536-43.