## **Rabbit Monoclonal Antibody to Human CD3**

| Catalog No.:           | RMPD 005   |
|------------------------|--|
| Intended Use:          | This product is intended for qualitative immunohistochemistry<br>with normal and neoplastic formalin-fixed, paraffin-embedded<br>tissue sections, to be viewed by light microscopy. Clinical<br>interpretation of staining results should be accompanied by<br>histological studies with proper controls. Patients' clinical<br>histories and other relevant diagnostic tests should be utilized by<br>a qualified person(s) when evaluating and interpreting results.   |
| Immunogen:             | A synthetic 13-mer peptide corresponding to aa 156-168 of the epsilon chain of human CD3 protein.  |
| Clone:                 | SP7  |
| Isotype:               | IgG  |
| Format:                | This antibody has been pretitered and quality controlled to work<br>on formalin-fixed paraffin-embedded and acetone fixed cryostat<br>tissue sections. No further titration is required.   |
| Staining Protocol:     | We suggest an incubation period of 30 minutes at room<br>temperature. Optimal incubation conditions should be<br>determined by the user based upon the fixation conditions and<br>staining system employed. Formalin fixed paraffin embedded<br>tissue sections require high temperature antigen unmasking with<br>10mM citrate buffer, pH 6.0 prior to immunostaining.  |
| Specificity:           | The CD3 antigen is present on early thymocytes and mature T cells and is generally regarded as a Pan T cell marker. This antibody will help detect CD3 expression in normal and neoplastic tissues. It reacts with the intracytoplasmic portion of the CD3 antigen expressed by T cells. It stains human T cells in both the cortex and medulla of the thymus and in peripheral lymphoid tissues. This antibody is suitable for staining normal and neoplastic T cells in formalin-fixed, paraffin-embedded tissues. It cross reacts with Human, Baboon, Monkey, Horse, Dog,, and Cat. |
| Positive Control:      | Tonsil   |
| Cellular Localization: | Cell Membrane  |
| Storage:               | Store at 2-8°C. Do not use beyond the expiration date stated on the label.   |

IVD: For In Vitro Diagnostic Use DBS will not be held responsible for patent infringement or other violation that may occur with the use of our product



DBS 1020 Serpentine Lane, # 114, Pleasanton, CA 94566 Tel: 925 484 3350, Fax: 925 484 3390 Website: <u>www.dbiosys.com</u> e-mail: customersupport@dbiosys.com