

# Goat Anti-Chicken IgY conjugated to SureLight® APC

 $\begin{array}{lll} \mbox{Product Number} & \mbox{D3-103} \\ \mbox{Amount} & \mbox{100 } \mbox{$\mu g$} \\ \mbox{Store at} & \mbox{2-8} \mbox{$^{\circ}$C} \end{array}$ 

## Form/Shipping & Storage

Supplied lyophilized. Upon receipt, store at -20  $^{\circ}$ C. Reconstitute with 1 ml of ddH2O and store at 2-8  $^{\circ}$ C. Phycobiliproteins are sensitive to freeze-thaw cycles.

## **Handling**

We recommend that the investigator determine the appropriate working concentration for their specific application. Avoid exposure to heat and light.

#### Buffer


Upon reconstitution, the product is in 10 mM tris (pH 8.2) + 100 mM NaCl + 0.01% sodium pentachlorophenol and 50mM Sucrose.

#### Stability

Product should be stored at 2-8°C in the dark and be used within 1 year. If further dilution of the conjugate is required, use diluted material within three months.

# <u>Note</u>

For research use only, not for diagnostic or therapeutic use.



Fluorescence excitation and emission spectra of APC in 100 mM sodium phosphate (pH 7.2) + 1 mM EDTA and 1 mM sodium azide. The emission scan was taken with excitation at 630 nm. The excitation scan was taken with an emission at 660 nm. The curves were normalized to equalize peak heights.

#### **Spectral Characteristics**

Visible absorption maxima 652 Emission maximum 657

## Concentration

Upon reconstitution with 1 ml ddH<sub>2</sub>O: 100µg/mL

Fluor:Protein ~1.4:1

