## **BioVision**

## **Calcein AM**


ALTERNATE NAME: Glycine,N,N'-[[3',6'-bis(acetyloxy)-3-oxospiro[isobenzofuran-1(3H),9'-[9H]xanthene]-2',7'-diyl]bis(methylene)]bis[N-[2-[(acetyloxy)methoxy]-2-oxoethyl]-,bis[(acetyloxy)methyl] ester; Calcein acetoxymethyl ester

CATALOG #:

1755-50, 250

AMOUNT:

STRUCTURE:



MOLECULAR FORMULA:	$C_{46}H_{46}N_2O_{23}$
MOLECULAR WEIGHT:	994.86
CAS NUMBER:	148504-34-1
APPEARANCE:	Off-white solid
SOLUBILITY:	DMSO (5 mg/ml)
PURITY:	≥95% by HPLC
STORAGE:	Store at -20 °C. Protect from light and moisture
DESCRIPTION:	A cell-permeable, non-fluorescent acetoxymethyl ester derivative of Calcein that gets hydrolyzed inside the cell by
	esterases. The loss of the acetoxymethyl group also enables calcein to readily bind intracellular calcium resulting in a strong yellowish-green fluorescence. It is the best probe for cell viability and cell adhesion due to its greater retention in cells as well and its pH insensitivity in physiological range.
HANDLING:	esterases. The loss of the acetoxymethyl group also enables calcein to readily bind intracellular calcium resulting in a strong yellowish-green fluorescence. It is the best probe for cell viablility and cell adhesion due to its greater retention in

FOR RESEARCH USE ONLY! Not to be used on humans.

## **RELATED PRODUCTS:**

- ApoSENSOR™ ATP Cell Viability Assay Kit (Cat. No. K254-100, 1000)
- VisionBlue™ Quick Cell Viability Assay Kit (Cat. No. K303-500, 2500)
- Live-Dead Cell Staining Kit (Cat. No.K501-100)
- Propidium Iodide (Cat. No. 1056-1)