

## Recombinant Human IL-27b/EBI-3 (Epstein-Barr Virus Induced Gene-3)

CATALOG #:	4376-20	20 µg
	4376-100	100 µg
	4376-1000	1 mg
LOT #:		
SOURCE:	<i>E. coli</i>	
PURITY:	> 90% by SDS-PAGE and RP-HPLC analysis Endotoxin level is < 0.01 ng per µg of rh- EBI-3.	
MOL. WEIGHT:	23.3 kDa	
FORM:	Lyophilized with mannitol.	

### RECONSTITUTION:

Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 – 0.5 mg/ml, which can be further diluted into other aqueous solutions.

### STORAGE CONDITIONS:

The lyophilized product is very stable at -20 °C. Upon reconstitution the material should be aliquoted and frozen at -20 °C. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

### DESCRIPTION:

Epstein-Barr Virus Induced Gene-3 (EBI-3), is a secreted glycoprotein belonging to the hematopoietin receptor family and related to the p40 subunit of IL-12. It was identified by its induced expression in B-lymphocytes in response to Epstein-Barr virus infection. EBI-3 forms heterodimers with p28 to form IL-27 and with p35 to form IL-35. Both IL-27 and IL-35 have anti-inflammatory and regulatory activity. Recombinant Human EBI is a non-glycosylated polypeptide chain consisting of 209 amino acids with a molecular weight of 23.3 kDa


### Biological Activity:

Assay data for human recombinant EBI-3 is based upon qualitative binding to anti-EBI-3 antibody.

### AA Sequence:

RKGPPAALTLPVQCRASRYPIAVDCSWTLPPAPNSTSPVSFIATYRLGMAARGHSWPCLQ  
QTPTSTSTCTITDVQLFSMAPYVLNVTAVHPWGSSSFVPFITEHIKPDPEGVRLSPLAERQL  
QVQWEPPGSPWPFPEIFSLKYWIRYKRQGAARFHRVGPIEATSFILRAVRPRARYYVQVAAQD  
LTDYGELSDWSLPATATMSLGK

**FOR RESEARCH USE ONLY! Not to be used in humans.**



HPLC analyses of IL-27. 20 µg of IL-27 were loaded on Phenomenex Gemini C18 column (250x4.6 mm 5 µm). The column was eluted with a 50 minute gradient of AcCN from 5 to 95% in 0.1% TFA at 1 ml/min.

### RELATED PRODUCTS:

#### Signal Transduction

- cAMP & cGMP Assay Kits
- Akt & JNK Activity Assay Kits
- Beta-Secretase Activity Assay Kit

#### Adipocyte & Lipid Transfer

- Recombinant Adiponectin, Survivin, & Leptin
- CETP Activity Assay & Drug Discovery Kits
- PLTP Activity Assay & Drug Discovery Kits
- Total Cholesterol Quantification Kit

#### Molecular Biology & Reporter Assays

- siRNA Vectors
- Cloning Insert Quick Screening Kit
- Mitochondrial & Genomic DNA Isolation Kits
- 5 Minutes DNA Ligation Kit
- 20 Minutes Gel Staining/Destaining Kit
- β -Galactosidase Staining Kit & Luciferase Reporter Assay Kit

#### Growth Factors and Cytokines

#### Monoclonal and Polyclonal Antibodies