

prosci-inc.com





## HIGH PERFORMANCE ANTIBODIES ... AND MORE


**ProSci Incorporated** 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

## **CHRM3 Antibody**

CATALOG NUMBER: 48-116

Background
BACKGROUND:



Immunohistochemistry staining of CHRM3 in brain cortex tissue using CHRM3 Antibody.

Specifications	
SPECIES REACTIVITY:	Chimpanzee, Gibbon, Gorilla, Human, Monkey, Orangutan
TESTED APPLICATIONS:	ELISA, IHC
APPLICATIONS:	CHRM3 antibody can be used in ELISA, and immunohistochemistry starting at 10 ug/mL.
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
SPECIFICITY:	BLAST analysis of the peptide immunogen showed no homology with other human proteins.
IMMUNOGEN:	CHRM3 antibody was raised against a peptide located in the 3rd cytoplasmic domain of CHRM3 (Human).
HOST SPECIES:	Rabbit
D "	
Properties	
PURIFICATION:	Immunoaffinity Chromatography
PHYSICAL STATE:	Liquid
BUFFER:	PBS, 0.1% sodium azide.
STORAGE CONDITIONS:	CHRM3 antibody should be stored long term (months) at -80 °C and short term (days) at 4 °C. As with all antibodies avoid freeze/thaw cycles.
CLONALITY:	Polyclonal
CONJUGATE:	Unconjugated
Additional Info	
ALTERNATE NAMES:	CHRM3, M3, Muscarinic receptor M3, Muscarinic M3 receptor, EGBRS, HM3, M3 muscarinic receptor
ACCESSION NO.:	P20309
PROTEIN GI NO.:	113125
OFFICIAL SYMBOL:	CHRM3
GENE ID:	1131

Muscarinic acetylcholine receptor 3 is an Acetylcholine Receptor that controls smooth muscle contraction by

mediating phosphoinositide hydrolysis and Ca2+ mobilization. Activation of this receptor causes secretion in glandular tissues. Mouse studies suggest that the M3 receptor may play a role in appetite stimulation and food intake.

## FOR RESEARCH USE ONLY

December 13, 2016