

prosci-inc.com




HIGH PERFORMANCE ANTIBODIES ... AND MORE

**ProSci Incorporated** 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

## **BACE1 Antibody**

CATALOG NUMBER: 48-373



Immunohistochemistry staining of BACE1 in pancreas tissue using BACE1 Antibody.

| Specifications       |  |
|----------------------|--|
| SPECIES REACTIVITY:  | Bat, Bovine, Dog, Gibbon, Guinea Pig, Hamster, Horse, Human, Monkey, Mouse, Pig, Rat   |
| TESTED APPLICATIONS: | IHC  |
| APPLICATIONS:        | BACE1 antibody can be used in immunohistochemistry starting at 1:50.   |
| USER NOTE:           | Optimal dilutions for each application to be determined by the researcher.   |
| SPECIFICITY:         | BLAST analysis of the peptide immunogen showed no homology with other human proteins.  |
| IMMUNOGEN:           | BACE1 antibody was raised against a peptide located near the internal domain of BACE1 (Human).   |
| HOST SPECIES:        | Rabbit   |
| <b>n</b> .:          |  |
| Properties           |  |
| PURIFICATION:        | Immunoaffinity Chromatography  |
| PHYSICAL STATE:      | Liquid   |
| BUFFER:              | PBS, 0.1% sodium azide.  |
| STORAGE CONDITIONS:  | BACE1 antibody should be stored long term (months) at -80 °C and short term (days) at 4 °C. As with all antibodies avoid freeze/thaw cycles.   |
| CLONALITY:           | Polyclonal   |
| CONJUGATE:           | Unconjugated   |
| Additional Info      |  |
| ALTERNATE NAMES:     | BACE1, APP beta-secretase, Asp 2, ASP2, Aspartyl protease 2, Aspartylprotease 2, Beta-secretase, Beta-site APP-cleaving enzyme, BACE, HSPC104, KIAA1149, B-Secretase, Beta-secretase 1, Memapsin-2 |
| ACCESSION NO .:      | P56817   |
| PROTEIN GI NO.:      | 296434407  |
| OFFICIAL SYMBOL:     | BACE1  |
| GENE ID:             | 23621  |
| Deekeround           |  |
| Background           |  |

BACKGROUND:

Bace 1 is responsible for the proteolytic processing of the amyloid precursor protein (APP). Cleaves at the N-

terminus of the A-beta peptide sequence, between residues 671 and 672 of APP, leads to the generation and extracellular release of beta-cleaved soluble APP, and a corresponding cell-associated C-terminal fragment which is later released by gamma-secretase.

FOR RESEARCH USE ONLY

December 13, 2016