

prosci-inc.com





## HIGH PERFORMANCE ANTIBODIES ... AND MORE


ProSci Incorporated 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

## **Cullin 1 Antibody**

CATALOG NUMBER: 49-412

Specifications



Immunohistochemistry staining of Cullin 1 in brain cortex tissue using Cullin 1 Antibody.

Specifications	
SPECIES REACTIVITY:	Human
TESTED APPLICATIONS:	ELISA, IHC, IP, WB
APPLICATIONS:	Cullin 1 antibody can be used in ELISA starting at 1:000 - 1:1000, and immunohistochemistry starting at 10 ug/mL.
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
IMMUNOGEN:	Cullin 1 antibody was raised against amino acids from the C-Terminus of Cullin 1 (Human).
HOST SPECIES:	Rabbit
Properties	
PURIFICATION:	Delipidation and Defibrination
PHYSICAL STATE:	Liquid
BUFFER:	0.01% sodium azide.
STORAGE CONDITIONS:	Store Cullin 1 antibody at 4 °C or -20 °C. As with all antibodies avoid freeze/thaw cycles.
CLONALITY:	Polyclonal
CONJUGATE:	Unconjugated
Additional Info	
ALTERNATE NAMES:	CUL1, Cullin 1, Cullin-1, CUL-1
ACCESSION NO.:	Q13616
PROTEIN GI NO.:	19863257
OFFICIAL SYMBOL:	CUL1
GENE ID:	8454

## **Background**

BACKGROUND:

Cullins assemble a potentially large number of ubiquitin ligases by binding to the RING protein ROC1 to catalyse polyubiquitination, as well as binding to various specificity factors to recruit substrates. Cullin 1 is an essential

component of the SCF (SKP1-CUL1-F-box protein) E3 ubiquitin ligase complex, which mediates the ubiquitination of proteins involved in cell cycle progression, signal transduction and transcription. In the SCF complex, cul1 serves as a rigid scaffold that organizes the SKP1-F-box protein and RBX1 subunits. Cul1 may also contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. Cul1 is part of the SCF complex consisting of CUL1, RBX1, SKP1 and SKP2, where it interacts directly with SKP1, SKP2 and RBX1. Cul1 also interacts with RNF7 and is part of a complex with TIP120A/CAND1 and RBX1. The unneddylated form interacts with TIP120A/CAND1 and the interaction negatively regulates the association with SKP1 in the SCF complex.

## FOR RESEARCH USE ONLY

December 13, 2016