

prosci-inc.com





HIGH PERFORMANCE ANTIBODIES ... AND MORE

ProSci Incorporated 12170 Flint Place Poway, CA 92064 Toll Free: +1 (888) 513 9525 Local: +1 (858) 513 2638 Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

CD178 Antibody [NOK-1]

CATALOG NUMBER: 76-936

Specifications	
SPECIES REACTIVITY:	Human
TESTED APPLICATIONS:	FACS, Func
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.
SPECIFICITY:	The NOK-1 monoclonal antibody specifically reacts with human CD178, which is the CD95 or Fas ligand.
HOST SPECIES:	Mouse
Properties	
PURIFICATION:	The monoclonal antibody was purified utilizing affinity chromatography. The endotoxin level is determined by LAL test to be less than 0.01 EU/ μ g of the protein.
PHYSICAL STATE:	liquid
BUFFER:	Phosphate-buffered aqueous solution, ph7.2.
CONCENTRATION:	2.0 mg/mL
STORAGE CONDITIONS:	The product should be stored undiluted at 4°C . Do not freeze.
CLONALITY:	Monoclonal
ISOTYPE:	Mouse IgG1, kappa
CONJUGATE:	Unconjugated
Additional Info	
ALTERNATE NAMES:	APTL, FASL, CD178, CD95L, ALPS1B, CD95-L, TNFSF6, APT1LG1, FASLG
OFFICIAL SYMBOL:	FASLG
GENE ID:	356
Background	
BACKGROUND:	The NOK-1 monoclonal antibody specifically reacts with human CD178, which is the CD95 or Fas ligand. CD178 is a TNF superfamily type II transmembrane glycoprotein expressed by activated T and NK cells and is involved in Fas-mediated apoptosis of lymphocytes. CD178 is also expressed by monocytes, neutrophils, granulocytes and the parenchymal cells of the retina and cornea. The NOK-1 antibody has been reported to bind to COOH-terminus of the Fas ligand in the region associated with Fas binding.
REFERENCES:	1) Kayagaki, N., Kawasaki, A., Ebata, T., Ohmoto, H., Ikeda, S., Inoue, S., Yagita, H. (1995). Metalloproteinase-mediated release of human Fas ligand. The Journal of experimental medicine, 182(6), 1777-1783.
	2) Oyaizu, N., Adachi, Y., Hashimoto, F., McCloskey, T. W., Hosaka, N., Kayagaki, N., Pahwa, S. (1997). Monocytes express Fas ligand upon CD4 cross-linking and induce CD4+ T cells apoptosis: a possible mechanism of bystander cell death in HIV infection. The Journal of Immunology, 158(5), 2456-2463.
	3) Villunger, A., Egle, A., Marschitz, I., Kos, M., Bck, G., Ludwig, H., Greil, R. (1997). Constitutive expression of Fas (Apo-1/CD95) ligand on multiple myeloma cells: a potential mechanism of tumor-induced suppression of immune surveillance.Blood,90(1), 12-20.