ProSci

prosci-inc.com



HIGH PERFORMANCE ANTIBODIES ... AND MORE

ProSci Incorporated
12170 Flint Place
Poway, CA 92064

Toll Free: +1 (888) 513 9525
Local: +1 (858) 513 2638
Fax: +1 (858) 513 2692

techsupport@prosci-inc.com

EpCAM Recombinant Protein

CATALOG NUMBER: 92-642

Specifications

SPECIES:	Human
SOURCE SPECIES:	Human Cells
SEQUENCE:	Gln24-Lys265
FUSION TAG:	C-Fc tag
APPLICATIONS:	This recombinant protein can be used for biological assays. For research use only.

Properties

PURITY:	Greater than 95% as determined by reducing SDS-PAGE. Endotoxin level less than 0.1 ng/ug (1 IEU/ug) as determined by LAL test.
PREDICTED MOLECULAR WEIGHT:	54.5 kD
PHYSICAL STATE:	Lyophilized
BUFFER:	Lyophilized from a 0.2 um filtered solution of PBS, pH7.4. It is not recommended to reconstitute to a concentration less than 100 ug/ml. Dissolve the lyophilized protein in ddH2O.
STORAGE CONDITIONS:	Lyophilized protein should be stored at -20˚C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7˚C for 2-7 days. Aliquots of reconstituted samples are stable at -20˚C for 3 months.

Additional Info

ALTERNATE NAMES:	Epithelial cell adhesion molecule, EpCAM, Trop-1, Epithelial cell surface antigen, EGP, hEGP314, TACSTD1, KS 1/4 antigen, KSA, CD326, GA733-2, M1S2, M4S1, MIC18, TROP1
ACCESSION NO.:	P16422

Background

Epithelial Cell Adhesion Molecule (EpCAM) is a signal type I transmembrane glycoprotein that belongs to the EPCAM family. EpCAM is composed of an extracellular domain with one thyroglobulin type-1 domain, a transmembrane domain and a cytoplasmic domain. EpCAM is found on the surface of adenocarcinoma, but not on mesodermal or neural cell membranes. The EpCAM molecule has been shown to function as a homophilic Ca2+ independent adhesion molecule. It may act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium as an immunological barrier providing the first line of defense against infection. Defects in EPCAM are a cause of hereditary non-polyposis colorectal cancer type 8 (HNPCC8) and diarrhea type 5 (DIAR5). EpCAM plays a role in embryonic stem cells proliferation and differentiation; it up-regulates the expression of FABP5, MYC and Cyclin A and Cyclin E. It is highly and selectively expressed by undifferentiated embryonic stem cells.

FOR RESEARCH USE ONLY

December 14, 2016